Dermatology Online Journal

Volume 14 Number 7

July 2008

[DOJ](#)

[Contents](#)

Localized abdominal idiopathic lipodystrophy

Vasco Vieira Serrão MD, Ana Barata Feio MD

Dermatology Online Journal 14 (7): 15

Abstract

Localized loss of subcutaneous tissue can occur after panniculitis, injections of corticosteroids and other drugs, or associated with infectious, autoimmune or neurologic diseases. The "idiopathic lipoatrophies" are a group of poorly characterized diseases, with focal disappearance of subcutaneous fat, and usually the thighs, abdomen or the ankles are affected. Three subtypes have been described based on clinical presentation: lipoatrophia semicircularis, annular lipoatrophy of the ankles and centrifugal lipodystrophy. We describe a 52-year-old female patient who developed a localized atrophy of the abdominal areas over a period of 3 months without any inflammatory signs over the evolution of the disease. The patient denied any previous local trauma or medication of any type. The atrophy stabilized, showing no progression over the last 6 years. The histopathological examination was normal except for the absence of subcutaneous fat, although the biopsy was taken down to the fascia. There was no clinical or serologic evidence of autoimmune diseases and laboratory testing for *Borrelia burgdorferi* infection was negative. Other causes of localized lipoatrophies were excluded and the final diagnosis was localized idiopathic lipodystrophy. Our patient is the second report on an abdominal lipodystrophy, with no previous inflammatory signs, absence of subcutaneous fat and no associated pathogenic factor. There is no established treatment for idiopathic lipodystrophy, and the lesions do not tend to resolve spontaneously.

Introduction

Localized loss of subcutaneous tissue is rare and may occur in a number of disease processes [1]. More frequently they represent

the resolution stage of a preceding panniculitis, usually associated with previous inflammatory signs [2]. Infectious, autoimmune and neurologic diseases may also be associated [2, 3, 4, 5] and a subgroup of patients will have no identifiable cause. These patients are grouped together as "localized idiopathic lipoatrophies," which represents a heterogeneous group of poorly characterized diseases [1, 6].

Case Report

Figure 1

Figure 1. Unilateral localized abdominal lipodystrophy, without previous inflammatory signs

Figure 2

Figure 2. Localized lipodystrophy, with complete loss of subcutaneous fat on the affected area

We describe a 52-year-old female patient, which developed a localized atrophy of the abdominal areas over a period of 3 months (Fig. 1). The subcutaneous tissue of the upper left abdominal quadrant progressively disappeared, leaving the so-called "pseudo-athletic" aspect, with notable veins and muscular tissue (Fig. 2). There were no surface skin changes, and the area was never indurated. The patient denied any inflammatory signs over the evolution of the disease, and there were no previous local trauma or medication of any type.

Figure 3

Figure 3. No changes in the affected area during the

The atrophy stabilized, showing no progression over the last 6 years (Fig. 3). The histopathological examination

evolution of the disease (pictures taken with an interval of two years)

had no epidermic or dermal pathological changes, with absence of subcutaneous fat although the biopsy was taken down to the fascia. An abdominal ultrasound and magnetic resonance of the area were performed, revealing only the atrophy of the subcutaneous fat, without associated muscular or organ changes. There was no clinical or serologic evidence of autoimmune diseases and laboratory testing for *Borrelia burgdorferi* infection was negative. Other causes of localized lipodystrophy were excluded and the final diagnosis was localized idiopathic lipodystrophy. Fat transfer was suggested, but the patient refused any treatment.

Discussion

In localized idiopathic lipoatrophy there is focal disappearance of subcutaneous fat, usually affecting the thighs, abdomen, or the ankles [2, 6]. Three subtypes have been described based on clinical presentation, with lipoatrophia semicircularis presenting as a band-like semicircular depression on the anterolateral aspects of the thighs probably due to repetitive trauma [6, 7, 8]. The presence of circumferential atrophic lesions of the ankles has also been described as annular lipoatrophy of the ankles, frequently associated with auto-immune diseases [9, 10]. Centrifugal lipodystrophy was initially described in Japanese children, with usually unilateral centrifugally spreading lipoatrophy, showing a bluish centre and erythema of the edges [11, 12]. There is only one previous report on an adult female with a clinical evolution similar to our patient, and was named unilateral localized idiopathic lipoatrophy [13]. In both patients localized unilateral loss of subcutaneous abdominal fat occurred, with no pre-existing inflammation or trauma. The overlying skin was normal and the lesions remained stable. In a review of panniculitis, Requena et al. defends that the designation of lipoatrophy should be used when there is a preceding inflammatory process involving the subcutis, corresponding to the end-stage of many types of panniculitis [2]. In contrast, lipodystrophy should be used when there is an absence of previous inflammatory stage and is more commonly associated with endocrinologic, metabolic, or autoimmune diseases [2]. Nevertheless, in the literature the 2 terms are often used interchangeably, adding further confusion to the nomenclature. As in both cases no previous clinical inflammation was described and histology could not identify any remaining subcutaneous tissue, the correct designation probably should be unilateral abdominal idiopathic lipodystrophy.

In the differential diagnosis of localized atrophies of the subcutaneous fat it must also be considered borreliosis and local injections of insulin, corticosteroids or other medication [3, 14, 15, 16, 17]. Panniculitis should also be excluded, where the presence of a preceding inflammatory stage is important [2]. The lack of epidermal or dermal changes differentiates localized idiopathic lipodystrophy from local panatrophy, which is a rare disorder with partial or total loss of subcutaneous fat, with associated atrophy of the overlying skin [18]. In panatrophy of Gowers, one of the two subtypes, single or multiple sharply defined areas of atrophy develop over a period of few weeks and in the affected area the subcutaneous tissue disappears [19]. There is no previous inflammatory process or sclerotic skin changes, and occurs mostly in young women [20]. Sclerotic panatrophy is another subtype, with sclerotic linear bands and occasional involvement of muscle or bone, occurring more frequently in children [21]. In both idiopathic lipoatrophy and panatrophy, after a rapid evolution, the lesions tend to remain unchanged indefinitely [13, 19]. Partial lipodystrophy is a more complex systemic disease, where there is disappearance of the fat from the face and upper half of the body, usually associated with progressive mesangiocapillary glomerulonephritis and hypocomplementemia [22]. There are rare cases of unilateral involvement, but the differential diagnosis is usually simple [22,23].

Two histopathological patterns have been described in localized lipoatrophies, with the involutional subtype being the most common, showing a pathological pattern with similarities to embryonic fat globules [24, 25]. In the inflammatory subtype there are normal lymphocytes, mixed inflammatory infiltrate and the presence of immunoreactants [26]. The biopsy in our patient was performed after the disease process had stabilized, and no subcutaneous fat was seen. The same happened in the other published case of unilateral localized idiopathic lipoatrophy and probably a biopsy taken in the early stages might have provided more information [13].

Although the pathogenesis is not known, there are associations of lipodystrophy to immunologically related diseases such as systemic lupus erythematosus, systemic sclerosis, autoimmune thyroid and renal disease [6, 23]. Repetitive trauma is considered the pathological factor involved in lipoatrophia semicircularis [7, 8]. It has also been suggested that there may be a primary or secondary abnormality of the sympathetic nervous system associated with localized disorders of the subcutaneous fat [26]. Lipodystrophy associated with antiretroviral treatment for the human immunodeficiency virus infection, has been related to changes of autonomic balance, resulting in redistribution of adipose tissue [27, 28]. The facial hemiatrophy of Parry-Romberg syndrome also seems to be related to a disorder of the sympathetic

nervous system [29].

There is no established treatment for idiopathic lipodystrophy and the lesions do not tend to resolve spontaneously [6, 13]. Autologous fat transfer to the atrophic site and the use of temporary or definitive fillers are therapeutical options [30, 31, 32, 33].

References

1. Garg A. Lipodystrophies. Am J Med. 2000 Feb;108(2):143-52. [PubMed](#).
2. Requena L, Sanchez Yus E. Panniculitis. Part II. Mostly lobular panniculitis. J Am Acad Dermatol. 2001 Sep;45(3):325-61. [PubMed](#).
3. Mullegger RR. Dermatological manifestations of Lyme borreliosis. Eur J Dermatol. 2004 Sep-Oct;14(5):296-309. [PubMed](#).
4. Quecedo E, Febrer I, Serrano G, Martinez-Aparicio A, Aliaga A. Partial lipodystrophy associated with juvenile dermatomyositis: report of two cases. Pediatr Dermatol. 1996 Nov-Dec;13(6):477-82. [PubMed](#).
5. Bilen N, Efendi H, Apaydin R, Bayramgurler D, Harova G, Komsuoglu S. Progressive facial hemiatrophy (Parry-Romberg syndrome). Australas J Dermatol. 1999 Nov;40(4):223-5. [PubMed](#).
6. Rongioletti F, Rebora A. Annular and semicircular lipoatrophy. Report of three cases and review of the literature. J Am Acad Dermatol. 1989 Mar;20(3):433-6. [PubMed](#)
7. Hodak E, David M, Sandbank M. Semicircular lipoatrophy - a pressure-induced lipoatrophy? Clin Exp Dermatol. 1990 Nov;15(6):464-5. [PubMed](#).
8. Herane MI, Urbina F, Sudy E. Lipoatrophia semicircularis: a compressive lipoatrophy consecutive to persistent mechanical pressure. J Dermatol. 2007 Jun;34(6):390-3. [PubMed](#).
9. Dimson OG, Esterly NB. Annular lipoatrophy of the ankles. J Am Acad Dermatol. 2006 Feb;54(2 Suppl):S40-2. [PubMed](#).
10. Roth DE, Schikler KN, Callen JP. Annular atrophic connective tissue panniculitis of the ankles. J Am Acad Dermatol. 1989 Nov;21(5 Pt 2):1152-6. [PubMed](#).
11. Hisamichi K, Suga Y, Hashimoto Y, Matsuba S, Mizoguchi M, Ogawa H. Two Japanese cases of localized involutional

lipoatrophy. *Int J Dermatol*. 2002 Mar;41(3):176-7. [PubMed](#).

12. Giam YC, Rajan VS, Hock OB. Lipodystrophia centrifugal abdominalis infantilis. *Br J Dermatol*. 1982 Apr;106(4):461-4. [PubMed](#).

13. Franks A, Verbov JL. Unilateral localized idiopathic lipoatrophy. *Clin Exp Dermatol*. 1993 Sep;18(5):468-9. [PubMed](#).

14. Levandoski LA, White NH, Santiago JV. Localized skin reactions to insulin: insulin lipodystrophies and skin reactions to pumped subcutaneous insulin therapy. *Diabetes Care*. 1982 May-Jun;5 Suppl 1:6-10. [PubMed](#).

15. Aviles-Izquierdo JA, Longo-Imedio MI, Hernanz-Hermosa JM, Lazaro-Ochaita P. Bilateral localized lipoatrophy secondary to a single intramuscular corticosteroid injection. *Dermatol Online J*. 2006 Mar 30;12(3):17. [PubMed](#).

16. Soos N, Shakery K, Mrowietz U. Localized panniculitis and subsequent lipoatrophy with subcutaneous glatiramer acetate (Copaxone) injection for the treatment of multiple sclerosis. *Am J Clin Dermatol*. 2004;5(5):357-9. [PubMed](#).

17. Kayikcioglu A, Akyurek M, Erk Y. Semicircular lipoatrophy after intragluteal injection of benzathine penicillin. *J Pediatr*. 1996 Jul;129(1):166-7. [PubMed](#).

18. Serup J, Weismann K, Kobayasi T, Ammitzboll T, Moller R. Local panatrophy with linear distribution: a clinical, ultrastructural and biochemical study. *Acta Derm Venereol*. 1982;62(2):101-5. [PubMed](#).

19. Sakamoto T, Oku T, Takigawa M. Gowers' local panatrophy. *Eur J Dermatol*. 1998 Mar;8(2):116-7. [PubMed](#).

20. Montagna P, Medori R, Zappia M, Gallassi R. Gowers local panatrophy. *Ital J Neurol Sci*. 1984 Dec;5(4):457-9. [PubMed](#).

21. Holmes RC, Meara RH. Morphoea, sclerotic panatrophy and disseminated granuloma annulare. *Clin Exp Dermatol*. 1983 Mar;8(2):201-3. [PubMed](#).

22. Levy Y, George J, Yona E, Shoenfeld Y. Partial lipodystrophy, mesangiocapillary glomerulonephritis, and complement dysregulation. An autoimmune phenomenon. *Immunol Res*. 1998 Aug;18(1):55-60. [PubMed](#).

23. Misra A, Peethambaram A, Garg A. Clinical features and metabolic and autoimmune derangements in acquired partial

lipodystrophy: report of 35 cases and review of the literature. *Medicine (Baltimore)*. 2004 Jan;83(1):18-34. [PubMed](#).

24. Peters MS, Winkelmann RK. The histopathology of localized lipoatrophy. *Br J Dermatol*. 1986 Jan;114(1):27-36. [PubMed](#).

25. Dahl PR, Zalla MJ, Winkelmann RK. Localized involutinal lipoatrophy: a clinicopathologic study of 16 patients. *J Am Acad Dermatol*. 1996 Oct;35(4):523-8. [PubMed](#).

26. Romijn JA, Fliers E. Sympathetic and parasympathetic innervation of adipose tissue: metabolic implications. *Curr Opin Clin Nutr Metab Care*. 2005 Jul;8(4):440-4. [PubMed](#).

27. Fliers E, Sauerwein HP, Romijn JA, Reiss P, van der Valk M, Kalsbeek A, Kreier F, Buijs RM. HIV-associated adipose redistribution syndrome as a selective autonomic neuropathy. *Lancet*. 2003 Nov 22;362(9397):1758-60. [PubMed](#).

28. van Gurp PJ, Tack CJ, van der Valk M, Reiss P, Lenders JW, Sweep FC, Sauerwein HP. Sympathetic nervous system function in HIV-associated adipose redistribution syndrome. *AIDS*. 2006 Mar 21;20(5):773-5. [PubMed](#).

29. Pinheiro TP, Silva CC, Silveira CS, Botelho PC, Pinheiro MG, Pinheiro Jde J. Progressive Hemifacial Atrophy--case report. *Med Oral Patol Oral Cir Bucal*. 2006 Mar 1;11(2):E112-4. [PubMed](#).

30. Butterwick KJ, Nootheti PK, Hsu JW, Goldman MP. Autologous fat transfer: an in-depth look at varying concepts and techniques. *Facial Plast Surg Clin North Am*. 2007 Feb;15(1):99-111. [PubMed](#).

31. Kaufman MR, Bradley JP, Dickinson B, Heller JB, Wasson K, O'Hara C, Huang C, Gabbay J, Ghadjar K, Miller TA. Autologous fat transfer national consensus survey: trends in techniques for harvest, preparation, and application, and perception of short- and long-term results. *Plast Reconstr Surg*. 2007 Jan;119(1):323-31. [PubMed](#).

32. Eppley BL, Dadvand B. Injectable soft-tissue fillers: clinical overview. *Plast Reconstr Surg*. 2006 Sep 15;118(4):98e-106e. [PubMed](#).

33. Broder KW, Cohen SR. An overview of permanent and semipermanent fillers. *Plast Reconstr Surg*. 2006 Sep;118(3 Suppl):7S-14S. [PubMed](#).